

DIFFERENTIAL FORMS AND THE WODZICKI RESIDUE.

WILLIAM J. UGALDE

ABSTRACT. For a pseudodifferential operator S of order 0 acting on sections of a vector bundle B on a compact manifold M without boundary, we associate a differential form of order dimension of M acting on $C^\infty(M) \times C^\infty(M)$. This differential form $\Omega_{n,S}$ is given in terms of the Wodzicki 1-density $\text{wres}([S, f][S, h])$. In the particular case of an even dimensional, compact, conformal manifold without boundary, we study this differential form for the case $(B, S) = (\mathcal{H}, F)$, that is, the Fredholm module associated by A. Connes [5] to the manifold M . We give its explicit expression in the flat case and then we address the general case. In [12] the author presented the computations in the six-dimensional case of a whole family of differential forms related to $\Omega_{n,S}$.

Keywords: Conformal geometry, Wodzicki residue, Fredholm module.

AMS Subject Classification: 53A30, 35S05, 58J40, 58B34, 46L87.

CONTENTS

1. Introduction	1
2. Properties of $\Omega_{n,S}$	3
2.1. Existence and uniqueness of $\Omega_{n,S}$	4
2.2. $\Omega_{n,S}$ and the Fredholm module on conformal manifolds	8
3. The leading symbol of F	8
4. Ω_n in the flat case	9
5. An approach to the general case	14
Appendix A. $\sigma(F)$ and the partial derivatives of the metric	17
References	20

1. INTRODUCTION

In [5], Connes associated to a even-dimensional compact conformal manifold M a canonical Fredholm module $(\mathcal{H}, F)$ and use it to define in the case $n = 4$, an n -dimensional differential form $\Omega_n(f, h)$ on M satisfying

$$(1) \quad \text{Wres}(f_0[F, f][F, h]) = \int_M f_0 \Omega_n(f, h)$$

for every f_0, f, h in $C^\infty(M)$. This differential form is determined by using the general formula for the total symbol of the product of two pseudodifferential operators, it happens to be *conformally invariant* and *uniquely* determined by the relation (1). In the 4-dimensional case, this differential form was found, by explicit

computations, to be *symmetric* and such that the Paneitz operator P relates to it by

$$\int_M \Omega_4(f, h) = 2 \int_M f P(h) dx.$$

Because of its potential relation with the study of conformally invariant differential operators generalizing the Yamabe operator, like the *GJMS operators* [8] for which the critical operator $P_4 = P/2$ in dimension $n = 4$ was recovered from Ω_4 , the study of such a differential form Ω_n , is of considerable importance in the case $n \geq 4$. In this paper we want to study this differential form in the following sense.

In Section 2.1, we associate to a pseudodifferential operator S of order 0, acting on sections of a vector bundle B of rank r on a compact manifold without boundary M , an n -dimensional differential form $\Omega_{n,S}$ acting on $C^\infty(M) \times C^\infty(M)$. $\Omega_{n,S}$ is obtained from the Wodzicki 1-density $\text{wres}([S, f][S, h])$ by using the formula for the total symbol of the product of two pseudodifferential operators (4). This differential form, will be uniquely determined by the relation

$$\text{Wres}(f_0[S, f][S, h]) = \int_M f_0 \Omega_{n,S}(f, h)$$

for every $f_0, f, h \in C^\infty(M)$. Furthermore $\int_M f_0 \Omega_{n,S}(f, h)$ define a Hochschild 2-cocycle on the algebra $C^\infty(M)$.

In Section 2.2 we will reduce ourselves to the particular case $(B, S) = (\mathcal{H}, F)$ of the Fredholm module associated by Connes [5] to an even-dimensional, compact, conformal manifold without boundary. We will prove that this differential form $\Omega_n = \Omega_{n,S}$ is indeed symmetric in f and h on any dimension. In Section 3 we compute the leading symbol of F , σ_L^F , and in Section 4, based on $\text{trace}(\sigma_L^F(\xi)\sigma_L^F(\eta))$ for ξ, η not zero in T_x^*M with the operator under the trace sign acting on middle dimension forms we will, in the flat case, express Ω_n as the integral of a specific polynomial obtained from the Taylor expansion of the function $\langle \xi, \eta \rangle^2 |\xi|^{-2} |\eta|^{-2}$ around a diagonal point. In Section 5, we write $\Omega_n(f, h)$ as $(Q_n(df, dh) + Q_{R,n}(df, dh))dx$ for a universal bilinear form which coincides with Ω_n in the flat case (see [5] for the case $n = 4$), and then we address the computations for the general case. At the end, we present an appendix on the dependence of the symbol of F on the partial derivatives of the metric.

In [12] the author presented a first glance at the computations of this differential form in the 6-dimensional case. The results on this paper are the rigorous proof of the statements needed for that computations, and any other done in general dimensions, to be true. My special thanks to T. Branson for unlimited support, and to J. C. Várilly for helpful comments on an earlier version.

Setting and Notations

Our general setting comes from [5] where, using the quantized calculus, Connes gives an analogue in dimension 4 of the 2-dimensional Polyakov action

$$I(X) = \frac{1}{2\pi} \int_\Sigma \eta_{ij} dX^i \wedge \star dX^j$$

for a Riemann surface Σ and a map X from Σ to $\mathbb{R}^d$. In the present work, unless otherwise stated, M represents an oriented, compact, manifold without boundary, endowed with a conformal structure, and of even dimension $n = 2l$ (which is going to be fixed from now on). $\mathcal{H}$ is the Hilbert space of square integrable forms of middle dimension, $\mathcal{H} = L^2(M, \Lambda_{\mathbb{C}}^l T^*M)$, on which functions on M act as multiplication

operators. F is the pseudodifferential operator of order 0 acting in $\mathcal{H}$, obtained from the orthogonal projection P on the image of d , by the relation $F = 2P - 1$. From the Hodge decomposition theorem [13] it is easy to see that F preserves the finite dimensional space of harmonic forms H^l , and that F restricted to the $\mathcal{H} \ominus H^l$ is given by

$$(2) \quad F = \frac{d\delta - \delta d}{d\delta + \delta d},$$

in terms of a Riemannian metric compatible with the conformal structure of M . Both $\mathcal{H}$ and F are independent of the metric in the conformal class [4, Section IV.4.γ].

The Riemann curvature tensor will be represented with the letter R , the Ricci tensor will be represented by $\text{Rc}_{ij} = R^k_{ikj}$, and the scalar curvature by $\text{Sc} = \text{Rc}^i_i$. If needed, we will “raise” and “lower” indices using the metric without explicit mention, for example, $g_{mi}R^i_{jkl} = R_{mjkl}$.

When working with the total symbol of a pseudodifferential operator P , we will denote its leading symbol by σ_L^P , or $\sigma_L(P)$ in case P has a long expression, and if the operator P is of order k then its total symbol (in some given local coordinates) will be represented as

$$\sigma(P) = \sigma_k^P + \sigma_{k-1}^P + \sigma_{k-2}^P + \cdots,$$

where $\sigma_L^P = \sigma_k^P$. It is important to note that the different σ_j^P for $j < k$ are defined only in local charts and are not diffeomorphism invariant [10]. However, Wodzicki [14] has shown that the term σ_{-n}^P enjoys a very special significance. For a pseudodifferential operator P , acting on sections of a vector bundle B over a manifold M , there is a 1-density on M expressed in local coordinates by

$$(3) \quad \text{wres}(P) = \int_{|\xi|=1} \{ \text{trace}(\sigma_{-n}^P(x, \xi)) d^{n-1}\xi \} dx.$$

This *Wodzicki residue density* is independent of the local representation. Here we are using the same notations as in [7], where an elementary proof of this matter can be found.

In the case of a Riemannian manifold (M, g) and an orientation dx , the metric g induces a metric g^1 on the cotangent bundle T^*M , and metrics g^k on the exterior bundles $\Lambda^k M$ in the following way [2]. If X is a vector field and ω_X denotes the 1-form determined by $\omega_X(Y) = g(X, Y)$ then $g^1(\omega_X, \omega_Y) := g(X, Y)$. If ω^i, η^j are 1-forms then,

$$g^k(\omega^1 \wedge \cdots \wedge \omega^k, \eta^1 \wedge \cdots \wedge \eta^k) := \det(g^1(\omega^i, \eta^j)).$$

Saying that dx is normalized means it has been chosen so that $g^n(dx, dx) = 1$.

2. PROPERTIES OF $\Omega_{n,S}$

For a pseudodifferential operator S of order 0 acting on sections of a vector bundle B over a compact manifold without boundary M , we consider P to be the pseudodifferential operator of order -2 given by the product $P = f_0[S, f][S, h]$ with each $f_0, f, h \in C^\infty(M)$. P is acting on the same vector bundle as S , where smooth functions on M act as multiplication operators. The total symbol of P , up to order $-n$, is represented as a sum of $r \times r$ matrices of the form

$$\sigma_{-2}^P + \sigma_{-3}^P + \cdots + \sigma_{-n}^P$$

where r is the rank of the vector bundle B . We aim to study

$$\text{Wres}(P) = \int_M \left\{ \int_{|\xi|=1} \text{trace}(\sigma_{-n}^P(x, \xi)) d^{n-1}\xi \right\} dx$$

where $\sigma_{-n}(x, \xi)$ is the component of order $-n$ in the total symbol of P , $|\xi| = 1$ means the Euclidean norm of the coordinate vector $(\xi_1, \dots, \xi_n)$ in $\mathbb{R}^n$, and $d^{n-1}\xi$ is the normalized volume on $\{|\xi| = 1\}$. $\text{Wres}(P)$ is independent of the choice of the local coordinates on M , the local basis of B , and defines a trace (see [14]).

2.1. Existence and uniqueness of $\Omega_{n,S}$. In general, the total symbol of the product of two pseudodifferential operators P_1 and P_2 is given by

$$(4) \quad \sigma(P_1 P_2) = \sum \frac{1}{\alpha!} \partial_\xi^\alpha (\sigma(P_1)) D_x^\alpha (\sigma(P_2))$$

where $\alpha = (\alpha_1, \dots, \alpha_n)$ is a multi-index, $\alpha! = \alpha_1! \cdots \alpha_n!$ and $D_x^\alpha = (-i)^{|\alpha|} \partial_x^\alpha$.

Lemma 2.1. *Let T be a pseudodifferential operator and for f in $C^\infty(M)$ let f represent the operator “multiplication by f ” then $\sigma(fT) = f\sigma(T)$ and in particular $\text{wres}(fT) = f \text{wres}(T)$, implying*

$$(5) \quad \text{wres}(f_0[S, f][S, h]) = f_0 \text{wres}([S, f][S, h]).$$

Proof. By (4) $\sigma(fT) = \sigma(f)\sigma(T)$ since $\sigma(f)$ does not depend on ξ , indeed $\sigma(f) = fI$ with I the identity operator on the sections the operator T is acting on, and the first assertion follows. In particular note that $\sigma_{-n}(fT) = f\sigma_{-n}(T)$.

By (3) we have

$$\begin{aligned} \text{wres}(fT) &= \left\{ \int_{|\xi|=1} \text{trace}(\sigma_{-n}(fT) d^{n-1}\xi) \right\} dx \\ &= \left\{ \int_{|\xi|=1} f \text{trace}(\sigma_{-n}(T) d^{n-1}\xi) \right\} dx = f \text{wres}(T), \end{aligned}$$

and the lemma follows by taking $T = [S, f][S, h]$. $\square$

Lemma 2.2. *$[S, f]$ is a pseudodifferential operator of order -1 with total symbol $\sigma([S, f]) = \sum_{k \geq 1} \sigma_{-k}([S, f])$ where*

$$(6) \quad \sigma_{-k}([S, f]) = \sum_{|\beta|=1}^k \frac{1}{\beta!} D_x^\beta(f) \partial_\xi^\beta (\sigma_{-(k-|\beta|)}^S).$$

Proof. Since S and f are of order zero, the commutator $[S, f]$ is of order minus one. The expansion of the symbol $\sigma([S, f])$ is given by

$$\begin{aligned}\sigma([S, f]) &= \sum_{|\beta| \geq 1} \frac{1}{\beta!} (\partial_\xi^\beta (\sigma^S) D_x^\beta (f)) \\ &= \sum_{|\beta|=1} \frac{1}{\beta!} D_x^\beta (f) (\partial_\xi^\beta (\sigma_0^S) + \partial_\xi^\beta (\sigma_{-1}^S) + \partial_\xi^\beta (\sigma_{-2}^S) + \cdots) \\ &\quad + \sum_{|\beta|=2} \frac{1}{\beta!} D_x^\beta (f) (\partial_\xi^\beta (\sigma_0^S) + \partial_\xi^\beta (\sigma_{-1}^S) + \cdots) \\ &\quad + \sum_{|\beta|=3} \frac{1}{\beta!} D_x^\beta (f) (\partial_\xi^\beta (\sigma_0^S) + \cdots) + \cdots \\ &= \sum_{k \geq 1} \left(\sum_{|\beta|=1}^k \frac{1}{\beta!} D_x^\beta (f) \partial_\xi^\beta (\sigma_{-(k-|\beta|)}^S) \right),\end{aligned}$$

and the result follows. $\square$

Lemma 2.3. *With the sum taken over $|\alpha'| + |\alpha''| + |\beta| + |\delta| + j + k = n$, $|\beta| \geq 1$, and $|\delta| \geq 1$,*

$$(7) \quad \begin{aligned}\sigma_{-n}([S, f][S, h]) \\ = \sum \frac{1}{\alpha'! \alpha''! \beta! \delta!} D_x^\beta (f) D_x^{\alpha'' + \delta} (h) \partial_\xi^{\alpha' + \alpha'' + \beta} (\sigma_{-j}^S) \partial_\xi^\delta (D_x^{\alpha'} (\sigma_{-k}^S)).\end{aligned}$$

Proof. By (4) we have

$$\sigma_{-n}([S, f][S, h]) = \sum_{r+s+|\alpha|=n} \frac{1}{\alpha!} \partial_\xi^\alpha (\sigma_{-r}([S, f])) D_x^\alpha (\sigma_{-s}([S, h])).$$

Using (6) and because $\partial_\xi^\alpha (D_x^\beta f) = 0$ for any $|\alpha| \geq 1$, we deduce

$$\begin{aligned}\sigma_{-n}([S, f][S, h]) &= \sum_{\substack{r+s+|\alpha|=n \\ r \geq 1, s \geq 1}} \frac{1}{\alpha!} \sum_{\substack{j+|\beta|=r \\ |\beta| \geq 1}} \frac{1}{\beta!} \partial_\xi^\alpha (D_x^\beta (f) \partial_\xi^\beta (\sigma_{-j}^S)) \sum_{\substack{k+|\delta|=s \\ |\delta| \geq 1}} \frac{1}{\delta!} D_x^\alpha (\partial_\xi^\delta (\sigma_{-k}^S) D_x^\delta (h)) \\ &= \sum_{\substack{r+s+|\alpha|=n \\ r \geq 1, s \geq 1}} \frac{1}{\alpha!} \sum_{\substack{j+|\beta|=r \\ |\beta| \geq 1}} \frac{1}{\beta!} D_x^\beta (f) \partial_\xi^{\alpha+\beta} (\sigma_{-j}^S) \times \\ &\quad \times \sum_{\substack{k+|\delta|=s \\ |\delta| \geq 1}} \frac{1}{\delta!} \sum_{\substack{0 \leq \alpha' \leq \alpha \\ \alpha' + \alpha'' = \alpha}} \frac{\alpha!}{\alpha'! \alpha''!} \partial_\xi^\delta (D_x^{\alpha'} (\sigma_{-k}^S)) D_x^{\alpha'' + \delta} (h),\end{aligned}$$

where we use, in the last equality, the Leibniz rule:

$$D_x^\alpha (fh) = \sum_{\substack{0 \leq \alpha' \leq \alpha \\ \alpha' + \alpha'' = \alpha}} \frac{\alpha!}{\alpha'! \alpha''!} D_x^{\alpha'} (f) D_x^{\alpha''} (h),$$

with α' a multi-index of the form $\alpha' = (\alpha'_1, \dots, \alpha'_n)$, and $\alpha' \leq \alpha$ meaning $\alpha'_i \leq \alpha_i$ for every $i = 1, \dots, n$, the sum taken over $|\alpha| + |\beta| + |\delta| + j + k = n$, $|\beta| \geq 1$, and $|\delta| \geq 1$. $\square$

By (5),

$$\begin{aligned}
 \text{wres}(f_0[S, f][S, h]) &= f_0 \left\{ \int_{|\xi|=1} \text{trace} \left\{ \sum \frac{1}{\alpha'! \alpha''! \beta! \delta!} D_x^\beta(f) D_x^{\alpha''+\delta}(h) \times \right. \right. \\
 &\quad \left. \left. \times \partial_\xi^{\alpha'+\alpha''+\beta}(\sigma_{-j}^S) \partial_\xi^\delta(D_x^{\alpha'}(\sigma_{-k}^S)) \right\} d^{n-1}\xi \right\} dx \\
 (8) \qquad \qquad \qquad &= f_0 \omega_{n,S}(f, h) dx = f_0 \Omega_{n,S}(f, h).
 \end{aligned}$$

which explicitly gives an expression for $\Omega_{n,S}(f, h)$.

Definition 2.4. For every $f, h \in C^\infty(M)$ we define by (8)

$$\Omega_{n,S}(f, h) = \omega_{n,S}(f, h) dx := \text{wres}([S, f][S, h]).$$

Because of the previous construction, we have the $\Omega_{n,S}$ is uniquely determined by its relation with the Wodzicki residue of the operator $f_0[S, f][S, h]$ as stated in the following theorem.

Theorem 2.5. There is a unique, n -differential form $\Omega_{n,S}$ such that

$$\text{Wres}(f_0[S, f][S, h]) = \int_M f_0 \Omega_{n,S}(f, h)$$

for all f_0, f, h in $C^\infty(M)$.

By construction, $\text{Wres}(f_0[S, f_1][S, f_2])$ is a Hochschild 2-cocycle over the algebra of smooth functions on a compact manifold, $\mathcal{A} = C^\infty(M)$. Next we want to state such a property. A Hochschild n -cochain φ on $\mathcal{A}$ is an $(n+1)$ -linear functional on $\mathcal{A}$. The coboundary operator, denoted by b , is given by:

$$\begin{aligned}
 (b\varphi)(f_0, \dots, f_{n+1}) &:= \\
 \sum_{j=0}^n (-1)^j \varphi(f_0, \dots, f_j f_{j+1}, \dots, f_{n+1}) &- (-1)^{n+1} \varphi(f_{n+1} f_0, \dots, f_n).
 \end{aligned}$$

The cohomology of this complex is the **Hochschild cohomology** of $\mathcal{A}$. In particular, a Hochschild 0-cocycle φ on $\mathcal{A}$ is a trace since $\varphi \in \mathcal{A}^* = \text{Hom}(\mathcal{A}, \mathbb{C})$, and $0 = (b\varphi)(f_0, f_1) = \varphi(f_0 f_1) - \varphi(f_1 f_0)$. A Hochschild 2-cocycle φ must satisfy

$$\begin{aligned}
 0 &= (b\varphi)(f_0, f_1, f_2, f_3) \\
 &= \varphi(f_0 f_1, f_2, f_3) - \varphi(f_0, f_1 f_2, f_3) + \varphi(f_0, f_1, f_2 f_3) - \varphi(f_3 f_0, f_1, f_2)
 \end{aligned}$$

for every $f_i \in C^\infty(M)$.

Lemma 2.6. $\text{Wres}(f_0[S, f_1][S, f_2])$ is a Hochschild 2-cocycle over the algebra of smooth functions on M .

Proof. Linearity is evident, and from the relation $[S, fh] = [S, f]h + f[S, h]$ we have, for $\varphi(f_0, f_1, f_2) = \text{Wres}(f_0[S, f_1][S, f_2])$:

$$\begin{aligned}
(b\varphi)(f_0, f_1, f_2, f_3) &= \text{Wres}(f_0 f_1 [S, f_2][S, f_3]) - \text{Wres}(f_0 [S, f_1 f_2][S, f_3]) \\
&\quad + \text{Wres}(f_0 [S, f_1][S, f_2 f_3]) - \text{Wres}(f_3 f_0 [S, f_1][S, f_2]) \\
&= \text{Wres}(f_0 f_1 [S, f_2][S, f_3]) \\
&\quad - \text{Wres}(f_0 [S, f_1] f_2 [S, f_3]) - \text{Wres}(f_0 f_1 [S, f_2][S, f_3]) \\
&\quad + \text{Wres}(f_0 [S, f_1][S, f_2] f_3) + \text{Wres}(f_0 [S, f_1] f_2 [S, f_3]) \\
&\quad - \text{Wres}(f_3 f_0 [S, f_1][S, f_2]) \\
&= \text{Wres}(f_0 [S, f_1][S, f_2] f_3) - \text{Wres}(f_3 f_0 [S, f_1][S, f_2]) = 0
\end{aligned}$$

because of the trace property of Wres . $\square$

So far, by taking $f_0 = 1$, by uniqueness, and by the trace property of Wres , we conclude

$$\int_M \Omega_{n,S}(f, h) = \text{Wres}([S, f][S, h]) = \text{Wres}([S, h][S, f]) = \int_M \Omega_{n,S}(h, f).$$

From (5), $\Omega_{n,S}(f, h) = \text{wres}([S, f][S, h])$ but in general wres is not a trace, hence asserting that $\Omega_{n,S}(f, h)$ is symmetric is asserting that

$$\text{wres}([S, f][S, h]) = \text{wres}([S, h][S, f]).$$

To conclude the symmetry of $\Omega_{n,S}(f, h)$ on f and h , it is necessary to request more properties on the operator S . As we will see, it is enough to have the property $S^2 f = f S^2$, for every $f \in C^\infty(M)$.

Theorem 2.7. *If $S^2 f = f S^2$ for every $f \in C^\infty(M)$ then the differential form $\Omega_{n,S}$ associated by Theorem 2.5 to (B, S) is symmetric in f and h .*

Proof. We are going to exploit the linearity and the trace property of the Wodzicki's residue [14]. Note that

$$\begin{aligned}
&\text{Wres}(f_0 [S, f][S, h] - f_0 [S, h][S, f]) \\
&= \text{Wres}(f_0 S f S h - f_0 S f h S - f_0 f S S h + f_0 f S h S \\
&\quad - f_0 S h S f + f_0 S h f S + f_0 h S S f - f_0 h S f S) \\
&= \text{Wres}(f_0 S f S h + f_0 f S h S - f_0 S h S f - f_0 h S f S)
\end{aligned}$$

using the fact S^2 commutes with any element of the algebra $C^\infty(M)$ and the commutativity of the algebra $C^\infty(M)$. By the trace property of the Wodzicki's residue, it follows that $\text{Wres}(f_0 F f F h) = \text{Wres}(h f_0 F f F)$ and $\text{Wres}(f_0 F h F f) = \text{Wres}(f f_0 F h F)$. Hence, by using once more the commutativity of $C^\infty(M)$ we conclude

$$\int_M f_0 \Omega_{n,S}(f, h) = \int_M f_0 \Omega_{n,S}(h, f)$$

for every $f_0, f, h \in C^\infty(M)$. Therefore $\Omega_{n,S}(f, h) = \Omega_{n,S}(h, f)$ because of the arbitrariness of f_0 . $\square$

2.2. $\Omega_{n,S}$ and the Fredholm module on conformal manifolds. For the rest of this paper we restrict ourselves to an even dimensional, compact, oriented, conformal manifold without boundary M , and (B, S) given by the canonical Fredholm module $(\mathcal{H}, F)$ associated to M by A. Connes [5]. In this particular case, the pseudodifferential operator of order 0 is given by $F = (d\delta - \delta d)(d\delta + \delta d)^{-1}$ acting on $\mathcal{H} = L^2(M, \Lambda_{\mathbb{C}}^l T^*M) \ominus H^l$, with H^l the finite dimensional space of middle dimension harmonic forms. By definition F is selfadjoint and such that $F^2 = 1$. We relax the notation by denoting $\Omega_n = \Omega_{n,F}$ in this particular situation.

Theorem 2.8. *In the particular case in which M is a even dimensional compact conformal manifold without boundary and $(\mathcal{H}, F)$ is the Fredholm module associated to M by A. Connes [5], there is a unique, symmetric, and conformally invariant n -differential form $\Omega_n = \Omega_{n,F}$ such that*

$$\text{Wres}(f_0[F, f_1][F, f_2]) = \int_M f_0 \Omega_n(f_1, f_2)$$

for all $f_i \in C^\infty(M)$. Furthermore, $\int_M f_0 \Omega_n(f_1, f_2)$ defines a Hochschild 2-cocycle on the algebra of smooth functions on M .

Proof. Uniqueness follows from (8). Symmetry follows from Theorem (2.7) and its conformal invariance follows from its construction. Indeed, as pointed out in [1, Section 2.1.2], $\text{Wres}(P)$ does not depend on the choice of the metric in the conformal class defining the cosphere bundle. The only possible metric dependence is given by the operator P . In our particular case $P = [F, f][F, h]$ is, as well as F , independent of the metric in the conformal class. Indeed, the only metric dependence of $(d\delta - \delta d)(d\delta + \delta d)^{-1}$ is the one given by the Hodge star operator $\star$ used in $\delta = -\star d\star$, which is invariant under conformal changes of the metric when restricted to middle dimension forms (see Lemma (2.9)). The last assertion follows from Lemma (2.6). $\square$

Lemma 2.9. *In the particular case of forms of middle dimension, the restriction $\star|_{\Lambda^l(T^*M)}$ is conformally invariant.*

Proof. If $\hat{g} = e^{2\eta}g$ is a conformally equivalent metric to the metric g , then on k -forms we have $\hat{g}^k = e^{-2k\eta}g^k$. In particular, with $\hat{d}x$ and $\hat{\star}$ the volume form and the Hodge star operator associated to the metric $\hat{g}$ we have

$$1 = \hat{g}^n(\hat{d}x, \hat{d}x) = e^{-2n\eta} g^n(\hat{d}x, \hat{d}x).$$

Now for two $(n-k)$ -forms ξ and ξ' with $\xi \wedge \star \xi' = g^{n-k}(\xi, \xi') dx$, we get

$$\xi \wedge \hat{\star} \xi' = \hat{g}^{n-k}(\xi, \xi') \hat{d}x = e^{-2(n-k)\eta} g^{n-k}(\xi, \xi') e^{n\eta} dx = e^{(2k-n)\eta} \xi \wedge \star \xi'.$$

and hence $\hat{\star} = e^{(2k-n)\eta} \star$. This shows that for $k = l$, the scaling has had no effect. $\square$

3. THE LEADING SYMBOL OF F

In this section we study the symbol of F in the flat case, and in general, its leading symbol. In [6, Lemma 1.5.3], it is proved that the leading symbols of d and δ are given by $\sigma_L(d) = i\varepsilon(\xi)$, $\sigma_L(\delta) = -i\iota(\xi)$, where $\xi \in T^*M$, and $\varepsilon(\xi)$ and $\iota(\xi)$ are the exterior and interior multiplications respectively. As a result of the identity $(i\varepsilon(\xi) - i\iota(\xi))^2 = |\xi|^2 I$ and since $\Delta = d\delta + \delta d = (d + \delta)^2$, in the same reference it is also proved that the leading symbol of Δ is given by $\sigma_L(\Delta) = |\xi|^2 I$.

Because of the rule for the leading symbol of the product of two pseudodifferential operators, $\sigma_L(P_1 P_2) = \sigma_L(P_1) \sigma_L(P_2)$, we conclude that $\sigma_L(\Delta^{-1}) = |\xi|^{-2} I$.

For the particular case of a flat metric, the total symbols of both $d\delta$ and δd coincide with their leading symbols, $\sigma(d\delta) = \sigma_L(d\delta)$ and $\sigma(\delta d) = \sigma_L(\delta d)$, which at the same time, do not depend on the variable x . As a consequence, the total symbols of both $\Delta = d\delta + \delta d$ and $d\delta - \delta d$ coincide with their leading symbols $\sigma(\Delta) = \sigma_L(\Delta)$ and $\sigma(d\delta - \delta d) = \sigma_L(d\delta - \delta d)$ in the flat case. It follows also that —see Lemma A.1— the total symbol of Δ^{-1} coincides with its leading symbol $\sigma(\Delta^{-1}) = \sigma_L(\Delta^{-1}) = |\xi|^{-2} I$ and furthermore, it does not depend on x , hence

$$\begin{aligned} \sigma(F)(x, \xi) &= \sigma(d\delta - \delta d)(x, \xi) \sigma(\Delta^{-1})(x, \xi) \\ &= |\xi|^{-2} (\varepsilon(\xi) \iota(\xi) - \iota(\xi) \varepsilon(\xi)). \end{aligned}$$

Summarizing, we have:

Proposition 3.1. *The leading symbol of F is given by*

$$\sigma_L(F)(x, \xi) = \sigma_L(F)(\xi) = |\xi|^{-2} (\varepsilon_\xi \iota_\xi - \iota_\xi \varepsilon_\xi)$$

for all $(x, \xi) \in T^*M$, $\xi \neq 0$. In the particular case of a flat metric, we also have $\sigma_{-k}(F) = 0$ for all $k \geq 1$.

4. Ω_n IN THE FLAT CASE

Let $\Omega_{n \text{ flat}}(f, h)$ be the n -dimensional form on a flat manifold M uniquely determined by Theorem 2.5. Since we are in the flat case we have $\sigma_{-k}^F(x, \xi) = 0$, for all $k > 0$. Because of that and (6),

$$\sigma_{-r}([F, f]) = \sum_{|\beta|=r} \frac{1}{\beta!} \partial_\xi^\beta (\sigma_L^F) D_x^\beta f.$$

Using this information we deduce from (7) with $\alpha' = 0$ and $\alpha'' = \alpha$,

$$\sigma_{-n}([F, f][F, h]) = \sum_{|\alpha|+|\beta|+|\delta|=n} \frac{1}{\alpha! \beta! \delta!} (D_x^\beta f) (D_x^{\alpha+\delta} h) (\partial_\xi^{\alpha+\beta} (\sigma_L^F)) (\partial_\xi^\delta (\sigma_L^F)),$$

with the sum taken over $|\alpha|+|\beta|+|\delta|=n$, $1 \leq |\beta|$, $1 \leq |\delta|$. By (8) and Theorem 2.5, we have:

Lemma 4.1.

$$\begin{aligned} \Omega_{n \text{ flat}}(f, h) &= \\ &\left\{ \int_{|\xi|=1} \text{trace} \left(\sum \frac{1}{\alpha! \beta! \delta!} (D_x^\beta f) (D_x^{\alpha+\delta} h) \partial_\xi^{\alpha+\beta} (\sigma_L^F) \partial_\xi^\delta (\sigma_L^F) \right) d^{n-1} \xi \right\} dx, \end{aligned}$$

with the sum taken over $|\alpha|+|\beta|+|\delta|=n$, $1 \leq |\beta|$, $1 \leq |\delta|$.

To better handle the previous expression, we consider

$$\phi(\xi, u, v) := \sum \frac{u^\beta v^{\alpha+\delta}}{\alpha! \beta! \delta!} \text{trace}(\partial_\xi^{\alpha+\beta} (\sigma_L^F) \partial_\xi^\delta (\sigma_L^F))$$

with the sum as before; in this way

$$\Omega_{n \text{ flat}}(f, h) = \left\{ \sum A_{a,b} (D_x^a f) (D_x^b h) \right\} dx$$

with

$$\sum A_{a,b} u^a v^b = \int_{|\xi|=1} \phi(\xi, u, v) d^{n-1} \xi.$$

Our next task is to compute the previous integral. Instead of a direct approach to compute $\text{trace}(\partial_\xi^{\alpha+\beta}(\sigma_L^F)\partial_\xi^\delta(\sigma_L^F))$, we shall try to use the Taylor expansion of the function

$$\psi(\xi, \eta) = \text{trace}(\sigma_L^F(\xi)\sigma_L^F(\eta)),$$

on the diagonal point (ξ, ξ) , as suggested in [5] for the 4-dimensional case. Indeed, with $\alpha = (\alpha_1, \dots, \alpha_n, \alpha_{n+1}, \dots, \alpha_{n+n}) = (\beta, \delta)$, we have

$$\begin{aligned} \psi(\xi + u, \eta + v) &= \sum_{|\alpha| \geq 0} \frac{(u, v)^\alpha}{\alpha!} \text{trace} \left(\partial_{(\xi, \eta)}^\alpha (\sigma_L^F(\xi)\sigma_L^F(\eta)) \Big|_{\eta=\xi} \right) \\ &= \sum_{|\beta|+|\delta| \geq 0} \frac{u^\beta v^\delta}{\beta! \delta!} \text{trace} \left(\partial_{(\xi, \eta)}^{(\beta, 0)} (\sigma_L^F(\xi)) \partial_{(\xi, \eta)}^{(0, \delta)} (\sigma_L^F(\eta)) \Big|_{\eta=\xi} \right) \\ &= \sum_{|\beta|+|\delta| \geq 0} \frac{u^\beta v^\delta}{\beta! \delta!} \text{trace} \left(\partial_\xi^\beta (\sigma_L^F(\xi)) \partial_\eta^\delta (\sigma_L^F(\eta)) \Big|_{\eta=\xi} \right). \end{aligned}$$

Therefore, the term of order n in the Taylor expansion of ψ at the point (ξ, ξ) is given by

$$\begin{aligned} T_n \psi(\xi, \xi, u, v) &= \sum_{|\beta|+|\delta|=n} \frac{u^\beta v^\delta}{\beta! \delta!} \text{trace} \left(\partial_\xi^\beta (\sigma_L^F(\xi)) \partial_\eta^\delta (\sigma_L^F(\eta)) \Big|_{\eta=\xi} \right) \\ &= \sum_{\substack{|\beta|+|\delta|=n \\ |\beta| \geq 1, |\delta| \geq 1}} \frac{u^\beta v^\delta}{\beta! \delta!} \text{trace} \left(\partial_\xi^\beta (\sigma_L^F(\xi)) \partial_\eta^\delta (\sigma_L^F(\eta)) \Big|_{\eta=\xi} \right) \\ &\quad + \sum_{|\beta|=n} \frac{u^\beta}{\beta!} \text{trace}(\partial^\beta (\sigma_L^F(\xi)) \sigma_L^F(\xi)) + \sum_{|\delta|=n} \frac{v^\delta}{\delta!} \text{trace} \left(\sigma_L^F(\xi) \partial_\eta^\delta (\sigma_L^F(\eta)) \Big|_{\eta=\xi} \right). \end{aligned}$$

We denote by $T'_n \psi(\xi, \eta, u, v)$ the term of order n in the Taylor expansion of $\psi(\xi, \eta)$ minus the terms with only powers of u or only powers of v . That is to say,

$$T'_n \psi(\xi, \eta, u, v) = \sum_{\substack{|\beta|+|\delta|=n \\ |\beta| \geq 1, |\delta| \geq 1}} \frac{u^\beta v^\delta}{\beta! \delta!} \text{trace}(\partial_\xi^\beta (\sigma_L^F(\xi)) \partial_\eta^\delta (\sigma_L^F(\eta))).$$

Now

$$\begin{aligned}
& T'_n \psi(\xi, \eta, u + v, v) \\
&= \sum_{\substack{|\beta|+|\delta|=n \\ |\beta| \geq 1, |\delta| \geq 1}} \frac{(u+v)^\beta v^\delta}{\beta! \delta!} \text{trace}(\partial_\xi^\beta(\sigma_L^F(\xi)) \partial_\eta^\delta(\sigma_L^F(\eta))) \\
&= \sum_{\substack{|\beta|+|\delta|=n \\ |\beta| \geq 1, |\delta| \geq 1}} \sum_{\beta'+\beta''=\beta} \frac{u^{\beta'} v^{\beta''+\delta}}{\beta'! \beta''! \delta!} \text{trace}(\partial_\xi^\beta(\sigma_L^F(\xi)) \partial_\eta^\delta(\sigma_L^F(\eta))) \\
&= \sum_{\substack{|\beta|+|\delta|=n \\ |\beta| \geq 1, |\delta| \geq 1}} \sum_{\substack{\beta'+\beta''=\beta \\ \beta' \neq 0}} \frac{u^{\beta'} v^{\beta''+\delta}}{\beta'! \beta''! \delta!} \text{trace}(\partial_\xi^\beta(\sigma_L^F(\xi)) \partial_\eta^\delta(\sigma_L^F(\eta))) \\
&\quad + \sum_{\substack{|\beta|+|\delta|=n \\ |\beta| \geq 1, |\delta| \geq 1}} \frac{v^{\beta+\delta}}{\beta! \delta!} \text{trace}(\partial_\xi^\beta(\sigma_L^F(\xi)) \partial_\eta^\delta(\sigma_L^F(\eta))) \\
&= \sum_{\substack{|\beta'|+|\beta''|+|\delta|=n \\ |\beta''| \geq 1, |\delta| \geq 1}} \frac{u^{\beta'} v^{\beta''+\delta}}{\beta'! \beta''! \delta!} \text{trace}(\partial_\xi^{\beta'+\beta''}(\sigma_L^F(\xi)) \partial_\eta^\delta(\sigma_L^F(\eta))) + T'_n \psi(\xi, \eta, v, v).
\end{aligned}$$

Therefore, by taking $\eta = \xi$ we obtain:

$$T'_n \psi(\xi, \xi, u + v, v) - T'_n \psi(\xi, \xi, v, v) = \phi(\xi, u, v),$$

which means that to compute ϕ we only need to compute the term of order n in the Taylor expansion of ψ at (ξ, ξ) and forget about the terms with only powers of u or v . Summarizing, we have

Theorem 4.2.

$$\Omega_n \text{flat}(f, h) = \left(\sum A_{a,b} (D_x^a f) (D_x^b h) \right) dx,$$

where

$$\sum A_{a,b} u^a v^b = \int_{|\xi|=1} (T'_n \psi(\xi, \xi, u + v, v) - T'_n \psi(\xi, \xi, v, v)) d^{n-1} \xi$$

and $T'_n \psi(\xi, \eta, u, v)$ is the term of order n in the Taylor expansion of the function $\psi(\xi, \eta) = \text{trace}(\sigma_L^F(\xi) \sigma_L^F(\eta))$ without the terms with only powers of u or only powers of v , at the point (ξ, η) .

A general formula for $\text{trace}(\sigma_L^F(\xi) \sigma_L^F(\eta))$

Because of Theorem (4.2), to obtain an explicit expression for Ω_n , at least for the flat case, it is necessary to study $\text{trace}(\sigma_L^F(\xi) \sigma_L^F(\eta))$ for ξ and η not zero in $T_x^* M$. Note that, ignoring the space of harmonic forms at each level, the operator $F = (d\delta - \delta d) \Delta^{-1}$, and its symbol, are defined and preserve the subspace of m -forms for any $0 \leq m \leq n$.

Theorem 4.3. With $\sigma_L^F(\xi) \sigma_L^F(\eta)$ acting on m -forms we have:

$$(9) \quad \text{trace}(\sigma_L^F(\xi) \sigma_L^F(\eta)) = a_{n,m} \frac{\langle \xi, \eta \rangle^2}{|\xi|^2 |\eta|^2} + b_{n,m},$$

where $\langle \xi, \eta \rangle$ represents the inner product $g^m(\xi, \eta)$ given by the metric and

$$\binom{n}{m} - a_{n,m} = b_{n,m} = \binom{n-2}{m-2} + \binom{n-2}{m} - 2 \binom{n-2}{m-1}.$$

To prove it, we need to set some notation and preliminary results. In the following ξ_i, η_i, ξ and η belong to T_x^*M and are not zero. Also we will represent the trace operator acting on the different subspaces with the same symbol, making clear out of the context on which dimension is it acting.

By using the trace identity $\text{trace}(AB) = \text{trace}(BA)$ and the relation

$$(10) \quad \varepsilon_{m-1}(\xi)\iota_m(\eta) + \iota_{m+1}(\eta)\varepsilon_m(\xi) = \langle \xi, \eta \rangle I_m$$

with I_m the identity on m -forms, we know, for every $m \geq 1$,

$$\text{trace}(\iota_m(\eta)\varepsilon_{m-1}(\xi)) + \text{trace}(\iota_{m+1}(\eta)\varepsilon_m(\xi)) = \binom{n}{m} \langle \xi, \eta \rangle.$$

In the particular case $m = 0$, we have $\iota_1(\eta)\varepsilon_0(\xi) = \langle \xi, \eta \rangle I_0$, so $\text{trace}(\iota_1(\eta)\varepsilon_0(\xi)) = \langle \xi, \eta \rangle$, and hence

Lemma 4.4. *For every $m \geq 0$,*

$$\begin{aligned} \text{trace}(\iota_{m+1}(\eta)\varepsilon_m(\xi)) &= \langle \xi, \eta \rangle \left(\binom{n}{m} - \binom{n}{m-1} + \dots + (-1)^m \binom{n}{0} \right) \\ &= A_{n,m} \langle \xi, \eta \rangle. \end{aligned}$$

By Proposition 3.1, $\sigma_L^F(x, \xi) = |\xi|^{-2}(\varepsilon_{m-1}(\xi)\iota_m(\xi) - \iota_{m+1}(\xi)\varepsilon_m(\xi))$, so

$$\begin{aligned} \text{trace}(\sigma_L^F(x, \xi)) &= |\xi|^{-2}(\text{trace}(\iota_m(\xi)\varepsilon_{m-1}(\xi)) - \text{trace}(\iota_{m+1}(\xi)\varepsilon_m(\xi))) \\ &= |\xi|^{-2} \left(\binom{n}{m} |\xi|^2 - 2 \text{trace}(\iota_{m+1}(\xi)\varepsilon_m(\xi)) \right). \end{aligned}$$

By (10),

$$(11) \quad \varepsilon_{m-1}(\xi)\iota_m(\xi) - \iota_{m+1}(\xi)\varepsilon_m(\xi) = 2\varepsilon_{m-1}(\xi)\iota_m(\xi) - |\xi|^2 I_m,$$

hence Proposition 3.1 implies

$$\begin{aligned} \text{trace}(\sigma_L^F(x, \xi)) &= |\xi|^{-2} \text{trace}(\varepsilon_{m-1}(\xi)\iota_m(\xi) - \iota_{m+1}(\xi)\varepsilon_m(\xi)) \\ &= 2|\xi|^{-2} \text{trace}(\varepsilon_{m-1}(\xi)\iota_m(\xi)) - \binom{n}{m} = 2A_{n,m-1} - \binom{n}{m}, \end{aligned}$$

and we obtain as a consequence of the previous Lemma:

Corollary 4.5. *For every $(x, \xi) \in T^*M$,*

$$\text{trace}(\sigma_L(F)(x, \xi)) = 2A_{n,m-1} - \binom{n}{m}.$$

Proof of Theorem 4.3. Using (11), and Lemma 4.4 we deduce

$$\begin{aligned} &\text{trace}(\sigma_L(F)(x, \xi)\sigma_L(F)(x, \eta)) \\ &= |\xi|^{-2}|\eta|^{-2} \text{trace}((2\varepsilon_{m-1}(\xi)\iota_m(\xi) - |\xi|^2 I_m)(2\varepsilon_{m-1}(\eta)\iota_m(\eta) - |\eta|^2 I_m)) \\ &= 4|\xi|^{-2}|\eta|^{-2} \text{trace}(\varepsilon_{m-1}(\xi)\iota_m(\xi)\varepsilon_{m-1}(\eta)\iota_m(\eta)) - 4A_{n,m-1} + \binom{n}{m}. \end{aligned}$$

Considering for any $m \geq 1$ the quantity

$$a_m(\xi_1, \xi_2, \eta_1, \eta_2) = \text{trace}(\varepsilon_{m-1}(\xi_1)\iota_m(\xi_2)\varepsilon_{m-1}(\eta_1)\iota_m(\eta_2)),$$

we have the relation

$$\begin{aligned} a_m(\xi_1, \xi_2, \eta_1, \eta_2) &= \text{trace}(((\xi_1, \xi_2)I_m - \iota_{m+1}(\xi_2)\varepsilon_m(\xi_1))(\langle \eta_1, \eta_2 \rangle I_m - \iota_{m+1}(\eta_2)\varepsilon_m(\eta_1))) \\ &= \langle \xi_1, \xi_2 \rangle \langle \eta_1, \eta_2 \rangle \left(\binom{n}{m} - 2A_{n,m} \right) + \text{trace}(\iota_{m+1}(\xi_2)\varepsilon_m(\xi_1)\iota_{m+1}(\eta_2)\varepsilon_m(\eta_1)), \end{aligned}$$

which implies

$$a_{m+1}(\eta_1, \xi_2, \xi_1, \eta_2) = a_m(\xi_1, \xi_2, \eta_1, \eta_2) + \langle \xi_1, \xi_2 \rangle \langle \eta_1, \eta_2 \rangle \left(2A_{n,m} - \binom{n}{m} \right),$$

where

$$\begin{aligned} a_1(\xi_1, \xi_2, \eta_1, \eta_2) &= \text{trace}(\varepsilon_0(\xi_1)\iota_1(\xi_2)\varepsilon_0(\eta_1)\iota_1(\eta_2)) \\ &= \text{trace}(\iota_1(\eta_2)\varepsilon_0(\xi_1)\iota_1(\xi_2)\varepsilon_0(\eta_1)) \\ &= \text{trace}(\langle \eta_2, \xi_1 \rangle \langle \xi_2, \eta_1 \rangle I_0) = \langle \eta_2, \xi_1 \rangle \langle \xi_2, \eta_1 \rangle. \end{aligned}$$

Thus, because of the relation

$$\text{trace}(\sigma_L(F)(x, \xi)\sigma_L(F)(x, \eta)) = 4a_m(\xi, \xi, \eta, \eta)|\xi|^{-2}|\eta|^{-2} + \binom{n}{m} - 4A_{n,m-1},$$

we have a recursive way of computing the left hand side. This is enough to prove that $\text{trace}(\sigma_L^F(\xi)\sigma_L^F(\eta)) = a_{n,m}\langle \xi, \eta^2 \rangle |\xi|^{-2}|\eta|^{-2} + b_{n,m}$ for some constants $a_{n,m}$ and $b_{n,m}$.

Now in the particular case in which $\xi = \eta$ is a member of an orthonormal basis of 1-forms, the operator under the trace equals I_m , the identity on m -forms because of the equality $\sigma_L(F)(\xi)\sigma_L(F)(\xi) = (\varepsilon(\xi)\iota(\xi) - \iota(\xi)\varepsilon(\xi))^2$ and the following computations. For a basic m -form $e_{j_1} \wedge \cdots \wedge e_{j_m}$, one has

$$(\varepsilon_{m-1}(\xi)\iota_m(\xi) - \iota_{m+1}(\xi)\varepsilon_m(\xi))(e_{j_1} \wedge \cdots \wedge e_{j_m}) = e_{j_1} \wedge \cdots \wedge e_{j_m}$$

if $\xi = e_{j_i}$ for some i ; and

$$(\varepsilon_{m-1}(\xi)\iota_m(\xi) - \iota_{m+1}(\xi)\varepsilon_m(\xi))(e_{j_1} \wedge \cdots \wedge e_{j_m}) = -e_{j_1} \wedge \cdots \wedge e_{j_m}$$

if $\xi \neq e_{j_i}$ for every i . Therefore $a_{n,m} + b_{n,m} = \binom{n}{m}$.

If both ξ and η are different members of an orthonormal basis, the term $\langle \xi, \eta \rangle$ vanishes and the expression (9) reduces to $b_{n,m}$, in this case,

$$\sigma_L(F)(\xi)\sigma_L(F)(\eta)e_{j_1} \wedge \cdots \wedge e_{j_m} = e_{j_1} \wedge \cdots \wedge e_{j_m}$$

if both $\xi, \eta \in \{e_{j_1}, \dots, e_{j_m}\}$ or if both $\xi, \eta \notin \{e_{j_1}, \dots, e_{j_m}\}$; and

$$\sigma_L(F)(\xi)\sigma_L(F)(\eta)e_{j_1} \wedge \cdots \wedge e_{j_m} = -e_{j_1} \wedge \cdots \wedge e_{j_m}$$

if only one of ξ, η belongs to $\{e_{j_1}, \dots, e_{j_m}\}$ and the other does not.

The number of basic m -forms containing both ξ and η as factors is $\binom{n-2}{m-2}$. The number of basic m -forms containing neither ξ nor η is $\binom{n-2}{m}$, and the number of basic m -forms containing exactly one of ξ or η as a factor is $2\binom{n-2}{m-1}$. In this way, the value of $b_{n,m}$ is given by the trace of the above operator, which equals $b_{n,m} = \binom{n-2}{m-2} + \binom{n-2}{m} - 2\binom{n-2}{m-1}$, and hence the value for $a_{n,m}$ is $a_{n,m} = \binom{n}{m} - \binom{n-2}{m-2} - \binom{n-2}{m} + 2\binom{n-2}{m-1}$, and the proof is complete. $\square$

We can now restate Theorem 4.2 as

Theorem 4.6.

$$\Omega_{n \text{ flat}}(f, h) = \left(\sum A_{a,b} (D_x^a f)(D_x^b h) \right) dx,$$

where

$$\sum A_{a,b} u^a v^b = \int_{|\xi|=1} (T'_n \psi(\xi, \xi, u+v, v) - T'_n \psi(\xi, \xi, v, v)) d^{n-1} \xi$$

and $T'_n \psi(\xi, \eta, u, v)$ is the term of order n in the Taylor expansion of the function $\psi(\xi, \eta) = a_{n,l} \langle \xi, \eta \rangle^2 |\xi|^{-2} |\eta|^{-2} + b_{n,l}$ without the terms with only powers of u or only powers of v , at the point (ξ, η) .

5. AN APPROACH TO THE GENERAL CASE

In this section we want to address a possible approach to the general case. This approach is the same used in [5] to compute Ω_4 and in [12] for Ω_6 . The first step to obtain Ω_n is to compute it explicitly in a flat metric. That is precisely the result given by Theorem 4.6. Then, by changing the metric conformally, one obtains the expression for Ω_n in the conformally flat case as follows. If g is a conformally flat metric, then $g = e^{2\eta} g_{\text{flat}}$ for a C^∞ function η and g_{flat} the flat metric on M . One expresses $\Omega_{n, \text{flat}}(f, h)$ in terms of the new metric g , and obtains an expression in terms of the covariant derivatives of η , f , h , and the Riemann curvature tensor R . By using the conformal change equation for the Ricci tensor:

$$(12) \quad \eta_{;ij} = -V_{ij} - \eta_{;i} \eta_{;j} + \frac{1}{2} \eta_{;k} \eta_{;k} g_{ij},$$

one replaces the second covariant derivatives on η with terms with the Ricci tensor. The result is the expression for $\Omega_n(f, h)$ in the conformally flat metric g .

In (12) V represents a normalized translation of the Ricci tensor, useful in conformal geometry, given in terms of the normalized scalar curvature J by

$$V = \frac{\text{Rc} - Jg}{n-2} \quad \text{with} \quad J = \frac{\text{Sc}}{2(n-1)},$$

and the indices after the semicolon represents covariant derivatives, $\eta_{;ij} = \nabla_j \nabla_i \eta$.

To study the conformal invariance, there is no need to study the whole conformal deformation. It is enough to study the conformal deformation up to order one in η as follows. If we set $\hat{g} = e^{2z\eta} g_{\text{flat}}$ where $\eta \in C^\infty(M)$ and z a constant, then the conformal variation of each expression is a polynomial in z whose coefficients are expressions in the metric and the conformal factor η (actually, this is an abuse of the language since the conformal factor is $e^{2z\eta}$). In this way, the conformal deformation up to order one in η is given by $\frac{d}{dz} \Big|_{z=0}$. If the conformal deformation of a natural tensor or a differential operator is zero up to order one then, by integration it is fully invariant, for details see [3].

The last step is to take the expression obtained in the conformally flat case and consider it in a general metric. Finally, study its conformal variation (up to order one), and then look for those terms that must be added to obtain a conformally invariant expression for a general conformally curved metric.

Following the spirit of [5], one reorder the summation of the expression given by (8) for $\Omega_n(f, h)$ by looking at $j+k+|\alpha'|$ in $\{0, 1, \dots, n-2\}$. In this way

$\text{trace}(\sigma_{-n}([F, f][F, h]))$ can be written as $\sum_{p=0}^{n-2} \text{trace}(\sigma_{-n}^{(p)})$, where

$$(13) \quad \text{trace}(\sigma_{-n}^{(p)}) = \sum \frac{1}{\alpha'! \alpha''! \beta! \delta!} D_x^\beta(f) D_x^{\alpha''+\delta}(h) \text{trace}(\partial_\xi^{\alpha'+\alpha''+\beta}(\sigma_{-j}^F) \partial_\xi^\delta(D_x^{\alpha'}(\sigma_{-k}^F))),$$

with the sum taken over $j+k+|\alpha'|=p$.

Each σ_{-j}^F is a $\binom{n}{l} \times \binom{n}{l}$ matrix with σ_0^F only invoking $g_{ij}(x)$ and the σ_{-j}^F for $j \geq 1$ are polynomial expressions on the partial derivatives of the metric at x , see Lemma (A.2).

By (13), $\text{trace}(\sigma_{-n}^{(0)})$ is given in terms of $g_{ij}(x)$ (without any of its partial derivatives) since only contains terms of the form $\text{trace}(\partial_\xi^{\alpha'+\alpha''+\beta}(\sigma_L^F) \partial_\xi^\delta(\sigma_L^F))$. Also by (13), any $\text{trace}(\sigma_{-n}^{(p)})$ for $p \geq 1$ is a polynomial expression on the partial derivatives of the metric at x with coefficients depending smoothly on $g_{ij}(x)$, indeed, each of these $\text{trace}(\sigma_{-n}^{(p)})$ only invoke terms of the form $\text{trace}(\partial_\xi^{\alpha'+\alpha''+\beta}(\sigma_{-j}^F) \partial_\xi^\delta(D_x^{\alpha'}(\sigma_{-k}^F)))$.

Evidently, these properties are preserved after integration over the variable ξ . By choosing the coordinates x^j to be geodesic normal coordinates at the point x , one can assume that $g_{ij}(x) = \delta_i^j$, that the first partial derivatives of the metric vanish at x , and that the higher partial derivatives are expressed in term of the Riemannian curvature and its covariant derivatives [9]. In this way, we have generalized Lemma 4 in [5] to any even dimension by splitting $\Omega_n(f, h)$ as the sum of

$$\int_{|\xi|=1} \text{trace}(\sigma_{-n}^{(0)}) d^{n-1}\xi dx = B_n(\nabla^a df, \nabla^b dh)$$

and

$$\left\{ \int_{|\xi|=1} \text{trace}(\sigma_{-n}^{(1)} + \dots + \sigma_{-n}^{(n-2)}) d^{n-1}\xi \right\} dx = C_n(R, df, dh).$$

In this way one has

Lemma 5.1. *There exists a universal bilinear expression $B_n(\nabla^\alpha df, \nabla^\beta dh)$ for some multi-indices α and β and an expression $C_n(R, df, dh)$ linear in f and h such that*

$$(14) \quad \Omega_n(f, h) = \{B_n(\nabla^\alpha df, \nabla^\beta dh) + C_n(R, df, dh)\} dx$$

where R is the Riemannian curvature tensor, ∇ the covariant differentiation, and $|\alpha| + |\beta| = n - 2$.

Note that covariant derivatives do not commute, making the notation $\nabla^\alpha df$ somehow ambiguous. It is assumed that $C(R, df, dh)$ will absorb any ambiguity. Also, because of the restriction $|\beta| \geq 1$, $|\delta| \geq 1$ in Lemma (2.3), $\Omega_n(f, h)$ depends only on df and dh rather than on f and h , that is why the right hand side of (14) is given in terms of df and dh .

In the particular case of a flat manifold, all the terms involving curvature vanish, in that situation $C_n(R, df, dh)$ reduces to zero and the covariant derivatives do commute. In particular, $B_n(\nabla^\alpha df, \nabla^\beta dh)$ equals $\Omega_n(f, h)$.

Lemma 5.2. *If M is a flat manifold,*

$$\Omega_n(f, h) = B_n(\nabla^\alpha df, \nabla^\beta dh) dx.$$

How to compute $C_n(R, df, dh)$? The expression $\omega_n(f, h)$ in (8) is a sum of homogeneous polynomials in the ingredients $\nabla^\alpha df$, $\nabla^\beta dh$, and $\nabla^\gamma R$ for multi-indices α, β , and γ , in the following sense, each monomial must satisfies the **homogeneity condition** given by the rule (see [3]):

$$2k_R + k_\nabla = n$$

where k_R denotes its degree in R and k_∇ its degree in ∇ . Also, for covariant derivatives we count all of the derivatives on R , f , and h , and any occurrence of Rc , or Sc as an occurrence of R . By closing under addition, we denote by $\mathcal{P}_n$ the space of these polynomials.

Because $|\beta| \geq 1$, and $|\delta| \geq 1$ in Lemma 2.3, we have $k_\nabla \geq 2$ and hence $k_R \leq (n-2)/2$. We say that Q is in $\mathcal{P}_{n,l}$ if Q can be written as a sum of monomials with $k_R \geq l$, or equivalently, $k_\nabla \leq n-2l$. We have

$$\mathcal{P}_n = \mathcal{P}_{n,0} \supseteq \mathcal{P}_{n,1} \supseteq \mathcal{P}_{n,2} \supseteq \cdots \supseteq \mathcal{P}_{n, \frac{n-2}{2}},$$

and $\mathcal{P}_{n,l} = 0$ for $l > (n-2)/2$.

There is an important observation to make. An expression which a priori appears to be in, say $\mathcal{P}_{6,1}$, may actually be in a subspace of it, like $\mathcal{P}_{6,2}$; for example,

$$\underbrace{f_{;i}h_{;jkl}W^{ijkl}}_{\in \mathcal{P}_{6,1}} = \underbrace{f_{;i}h_{;j}V_{kl}W^{ikjl} + f_{;i}h_{;j}W^i{}_{klm}W^j{}_{klm}}_{\in \mathcal{P}_{6,2}},$$

by reordering covariant derivatives and making use of the symmetries of the Weyl tensor W . Because of this filtration, it is necessary to adopt a fixed convention on how the indices should be placed when representing each expression in its index notation. For example, $f_{;ijk}h^{ijk}$ will be preferred over $f_{;ikj}h^{ijk}$ or any other variation. Lemma (5.1) is restated as

Lemma 5.3. *There exists a universal bilinear form $Q_n(df, dh)$ in $\mathcal{P}_{n,0} \setminus \mathcal{P}_{n,1}$ and a form $Q_{R,n}(df, dh)$ in $\mathcal{P}_{n,1}$ such that*

$$\omega_n(f, h) = Q_n(df, dh) + Q_{R,n}(df, dh)$$

where R is the Riemannian curvature tensor.

Once one has computed the expression for Ω_n in a flat metric g , one express it in a conformally related metric $\hat{g}$. By doing that, all of the terms for $Q_{R,n}$ which do not involve a factor of the Weyl tensor W are computed. In this way, up to the conformally flat case, one obtains the expression for Ω_n . Next, one uses the homogeneity condition $2k_\nabla + k_R = n$ to list the possible terms to be considered in the general conformally curved case. These terms must be such that they contain the Weyl tensor as a factor, and their conformal variation compensate that of the already known terms for Ω_n .

In the particular case $n = 4$, as found in [5], the homogeneity condition shows that no extra terms exist and the expression found in the conformally flat case suffices. Indeed, $\Omega_4(f, h)$ is given by a constant multiple of

$$-f_{;ij}{}^jh_{;i}{}^i - \frac{1}{2}f_{;i}{}^ih_{;j}{}^j - f_{;ij}h^{ij} - f_{;i}h^i{}_{;j}{}^j + 2f_{;i}h^i{}_{;j}J,$$

where $2(n-1)J = Sc$. The only terms to be added, $f_{;i}h^i{}_{;j}{}^j W^{***}$ or $f_{;i}h_{;j}W^{i*j*}$, are zero because of the symmetries of the Weyl tensor W (here each $*$ represents an index).

In the case $n = 6$, as in [12, Section 6], six extra terms are needed as a result of the homogeneity condition. As shown in that reference, symmetry, conformal invariance, and even the Hochschild 2-cocycle property of $\int_M f_0 \Omega_n(f_1, f_2)$ are not enough to compute the coefficients in front of those extra terms containing the Weyl tensor as a factor, indeed, it is not possible to determine, out of those three properties, the coefficients for $f_{;i}h_{;j}W_{jklm}W^{jklm}$ and $f_{;i}h_{;j}W^i{}_{klm}W^{jklm}$.

APPENDIX A. $\sigma(F)$ AND THE PARTIAL DERIVATIVES OF THE METRIC

One of the basis of the result of [5] is its Lemma 3, where M is a 4-dimensional manifold and the operator F , as well as its symbol, are acting on 2-forms. Our objective in this paragraph is to give a more general version of this lemma, in which we try to understand the behavior up to any sub-index, of the component σ_{-k} of the total symbol of F . At the same time, we give a detailed computation of the total symbol of Δ^{-1} in terms of the total symbol of Δ , and an explicit recursive computation of the total symbol of F in terms of the total symbols of Δ^{-1} and of $d\delta - \delta d$ in some given local coordinates.

To simplify the typing, we will abbreviate using the notation $T \sim \mathcal{L}_k$ to mean “the term T has the property of being linear in the k -partial derivatives of the metric at x with coefficients depending smoothly on the $g_{ij}(x)$ ”. By $T \sim (\mathcal{L}_k)^r$ we mean “the term T has the property of being a sum of products of r linear expressions of the type $\mathcal{L}_k$ ”. By $T \sim \mathcal{L}_0$ we mean “the term T only invokes $g_{ij}(x)$ ”, thus if $T \sim (\mathcal{L}_0)(\mathcal{L}_k)$ then $T \sim \mathcal{L}_k$.

The first step is an recursive computation of the symbol of Δ^{-1} in terms of the symbol of Δ . We use the following reasoning, justified by [11, Corollary 1.4.3]. From [6, Lemma 2.4.4], the operator $\Delta = d\delta + \delta d$ acting on $\Lambda^l T^*M$ can be expressed in a given system of local coordinates in the form $\sigma(d\delta + \delta d) = p_2 + p_1 + p_0$, where each p_i is a $\binom{n}{l} \times \binom{n}{l}$ matrix such that $p_2 \sim \mathcal{L}_0$, $p_1 \sim \mathcal{L}_1$, and $p_0 \sim \mathcal{L}_2 + (\mathcal{L}_1)^2$.

Let

$$\sigma(\Delta^{-1}) = \sum_{k \leq -2} \sigma_k(\Delta^{-1}) = \sum_{k \leq -2} r_k$$

be the total symbol of Δ^{-1} , hence

Lemma A.1. *The total symbol of Δ^{-1} is given by the recursive formula $r_{-2} = p_2^{-1}$, $r_{-3} = p_2^{-1}(-p_1 p_2^{-1} + \partial_\xi^1 p_2 D_x^1(p_2^{-1}))$, and in general, for all $k \geq 2$,*

$$\begin{aligned} r_{-(k+2)} = & p_2^{-1}(-p_1 r_{-(k+1)} - p_0 r_{-k} \\ & - (\partial_\xi^1 p_2 D_x^1 r_{-(k+1)} + \partial_\xi^1 p_1 D_x^1 r_{-k}) - \frac{1}{2} \partial_\xi^2 p_2 D_x^2 r_{-k}). \end{aligned}$$

Proof.

$$\begin{aligned}
(15) \quad 1 &= \sum_{\alpha} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} \sigma(\Delta) D_x^{\alpha} \sigma(\Delta^{-1}) = \sum_{|\alpha|=0}^2 \frac{1}{\alpha!} \partial_{\xi}^{\alpha} \sigma(\Delta) D_x^{\alpha} \sigma(\Delta^{-1}) \\
&= (p_2 + p_1 + p_0)(r_{-2} + r_{-3} + r_{-4} + \cdots) \\
&\quad + (\partial_{\xi}^1 p_2 + \partial_{\xi}^1 p_1)(D_x^1 r_{-2} + D_x^1 r_{-3} + D_x^1 r_{-4} + \cdots) \\
&\quad + \frac{1}{2}(\partial_{\xi}^2 p_2)(D_x^2 r_{-2} + D_x^2 r_{-3} + D_x^2 r_{-4} + \cdots) \\
&= [p_2 r_{-2}] + [p_2 r_{-3} + p_1 r_{-2} + \partial_{\xi}^1 p_2 D_x^1 r_{-2}] \\
&\quad + \sum_{k=2} [p_2 r_{-(k+2)} + p_1 r_{-(k+1)} + p_0 r_{-k} \\
&\quad + (\partial_{\xi}^1 p_2 D_x^1 r_{-(k+1)} + \partial_{\xi}^1 p_1 D_x^1 r_{-k}) + \frac{1}{2} \partial_{\xi}^2 p_2 D_x^2 r_{-k}] + \cdots
\end{aligned}$$

where $D_{\beta}^m = \sum_{|\alpha|=m} D_{\beta}^{\alpha}$ and each term in the summation has homogeneous order k in the variable ξ . The lemma follows by solving for r_{-k} at each level of the decomposition. $\square$

Lemma A.2. *Acting on middle dimension forms, the total symbol of Δ^{-1} is given as a sum of $\binom{n}{l} \times \binom{n}{l}$ matrices of the form $\sigma(\Delta^{-1}) = r_{-2} + r_{-3} + r_{-4} + \cdots$ where $r_{-2} \sim \mathcal{L}_0$, $r_{-3} \sim \mathcal{L}_1$, $r_{-4} \sim \mathcal{L}_2 + (\mathcal{L}_1)^2$, and in general, for $k \geq 0$,*

$$r_{-(k+2)} \sim \sum (\mathcal{L}_k)^{a_k} (\mathcal{L}_{k-1})^{a_{k-1}} \cdots (\mathcal{L}_1)^{a_1}$$

with the summation taken over $0 \leq a_i \leq k$, for every i and $a_1 + 2a_2 + \cdots + ka_k = k$.

Proof. The differentiation ∂_{ξ}^{α} does not alter the properties of p_k , so that, for instance, $\partial_{\xi}^{\alpha} p_1 \sim \mathcal{L}_1$. The differentiation D_x^{α} behaves in the following way $D_x^{\alpha}(\mathcal{L}_k) \sim \mathcal{L}_{k+|\alpha|}$ and $D_x(\mathcal{L}_k \mathcal{L}_j) \sim \mathcal{L}_{k+1} \mathcal{L}_j + \mathcal{L}_k \mathcal{L}_{j+1}$. The lemma follows from the next inductive reasoning:

The first three cases read directly from the expressions for r_{-j} in Lemma A.1, with $j = 2, 3, 4$. For the general case, assume

$$\begin{aligned}
(16) \quad r_{-(k+2)} &\sim \mathcal{L}_1 r_{-(k+1)} + \mathcal{L}_2 r_{-k} D_x^1 r_{-(k+1)} + \mathcal{L}_1 D_x^1 r_{-k} + D_x^2 r_{-k} \\
&\sim \mathcal{L}_1 \sum (\mathcal{L}_{k-1})^{a_{k-1}} \cdots (\mathcal{L}_1)^{a_1} + \mathcal{L}_2 \sum (\mathcal{L}_{k-2})^{b_{k-2}} \cdots (\mathcal{L}_1)^{b_1} \\
&\quad + D_x^1 \sum (\mathcal{L}_{k-1})^{a_{k-1}} \cdots (\mathcal{L}_1)^{a_1} + \mathcal{L}_1 D_x^1 \sum (\mathcal{L}_{k-2})^{b_{k-2}} \cdots (\mathcal{L}_1)^{b_1} \\
&\quad + D_x^2 \sum (\mathcal{L}_{k-2})^{b_{k-2}} \cdots (\mathcal{L}_1)^{b_1}
\end{aligned}$$

where $a_1 + 2a_2 + \cdots + (k-1)a_{k-1} = k-1$ and $b_1 + 2b_2 + \cdots + (k-2)b_{k-2} = k-2$.

First note that

$$\mathcal{L}_1 \sum (\mathcal{L}_{k-1})^{a_{k-1}} \cdots (\mathcal{L}_1)^{a_1} = \sum (\mathcal{L}_{k-1})^{a_{k-1}} \cdots (\mathcal{L}_1)^{1+a_1}$$

where $(1 + a_1) + 2a_2 + \cdots + (k-1)a_{k-1} = 1 + k - 1 = k$, and

$$\mathcal{L}_2 \sum (\mathcal{L}_{k-2})^{b_{k-2}} \cdots (\mathcal{L}_1)^{b_1} = \sum (\mathcal{L}_{k-2})^{b_{k-2}} \cdots (\mathcal{L}_2)^{1+b_2} (\mathcal{L}_1)^{b_1}$$

where $b_1 + 2(1 + b_2) + \cdots + (k-2)b_{k-2} = 2 + k - 2 = k$. In this way, the first two terms of (16) have the required property.

Next,

$$\begin{aligned}
& D_x^1 \sum (\mathcal{L}_{k-1})^{a_{k-1}} \cdots (\mathcal{L}_1)^{a_1} \\
&= \sum \mathcal{L}_k (\mathcal{L}_{k-1})^{a_{k-1}} \cdots (\mathcal{L}_1)^{a_1} + \sum (\mathcal{L}_{k-1})^{a_{k-1}+1} (\mathcal{L}_{k-2})^{a_{k-2}-1} \cdots (\mathcal{L}_1)^{a_1} + \cdots \\
(17) \quad & + \sum (\mathcal{L}_{k-1})^{a_{k-1}} \cdots (\mathcal{L}_2)^{a_2+1} (\mathcal{L}_1)^{a_1-1},
\end{aligned}$$

where each of the summations is taken over a set of the form $(d_1, \dots, d_k)$ with $d_1 + 2d_2 + \cdots + kd_k = k$. Indeed,

$$a_1 + \cdots + (k-1)(a_{k-1}-1) + k = k-1 - (k-1) + k = k,$$

$$a_1 + \cdots + (k-2)(a_{k-2}-1) + (k-1)(a_{k-1}+1) = k-1 - (k-2) + (k-1) = k,$$

and

$$(a_1-1) + 2(a_2+1) + \cdots + (k-1)a_{k-1} = 1 + k-1 = k.$$

The previous reasoning shows that $D_x^1 \left(\sum (\mathcal{L}_{k-2})^{b_{k-2}} \cdots (\mathcal{L}_1)^{b_1} \right)$ is a sum over a set of indices of the form $(a_1, \dots, a_{k-1})$ with $a_1 + 2a_2 + \cdots + (k-1)a_{k-1} = k-1$, thus

$$\mathcal{L}_1 D_x^1 \sum (\mathcal{L}_{k-2})^{b_{k-2}} \cdots (\mathcal{L}_1)^{b_1} \sim \mathcal{L}_1 \sum (\mathcal{L}_{k-1})^{a_{k-1}} \cdots (\mathcal{L}_1)^{a_1},$$

which satisfies the desired property as before. For the last term in (16),

$$D_x^2 \left(\sum (\mathcal{L}_{k-2})^{b_{k-2}} \cdots (\mathcal{L}_1)^{b_1} \right) \sim D_x \left(D_x \left(\sum (\mathcal{L}_{k-2})^{b_{k-2}} \cdots (\mathcal{L}_1)^{b_1} \right) \right),$$

which is of the desired type by applying twice the reasoning on (17). $\square$

Lemma A.3. *Acting on middle dimension forms, the total symbol σ^F of F , up to order $-k$ inclusive, is a $\binom{n}{l} \times \binom{n}{l}$ matrix of the form $\sigma^F = \sigma_0^F + \sigma_{-1}^F + \cdots + \sigma_{-k}^F + \cdots$ where $\sigma_0^F \sim \mathcal{L}_0$, $\sigma_{-1}^F \sim \mathcal{L}_1$, $\sigma_{-2}^F \sim \mathcal{L}_2 + (\mathcal{L}_1)^2$, and in general*

$$\sigma_{-k}^F \sim \sum (\mathcal{L}_k)^{\alpha_k} (\mathcal{L}_{k-1})^{\alpha_{k-1}} \cdots (\mathcal{L}_1)^{\alpha_1}$$

with the summation taken over $0 \leq \alpha_i \leq k$, for every i and $\alpha_1 + 2\alpha_2 + \cdots + k\alpha_k = k$.

Proof. The operator $d\delta - \delta d$ satisfies $\sigma(d\delta - \delta d) = q_2 + q_1 + q_0$, with q_2 of type $\mathcal{L}_0$, q_1 of type $\mathcal{L}_1$, and q_0 of type $\mathcal{L}_2 + (\mathcal{L}_1)^2$; these results follow exactly as the result for $\Delta = d\delta + \delta d$.

As in (15) we have, for the symbol of $F = (d\delta - \delta d)\Delta^{-1}$, the expansion

$$\begin{aligned}
\sigma(F) &= [q_2 r_{-2}] + [q_2 r_{-3} + q_1 r_{-2} + \partial_\xi^1 q_2 D_x^1 r_{-2}] \\
&+ \sum_{k=2} [q_2 r_{-(k+2)} + q_1 r_{-(k+1)} + q_0 r_{-k} \\
&+ (\partial_\xi^1 q_2 D_x^1 r_{-(k+1)} + \partial_\xi^1 q_1 D_x^1 r_{-k}) + \frac{1}{2} \partial_\xi^2 q_2 D_x^2 r_{-k}] + \cdots \\
&= \sigma_0^F + \sigma_{-1}^F + \sigma_{-2}^F + \cdots.
\end{aligned}$$

From this last equality and the properties of q_i and r_j , we get that $\sigma_0^F \sim \mathcal{L}_0$, $\sigma_{-1}^F \sim \mathcal{L}_1$, $\sigma_{-2}^F \sim \mathcal{L}_2 + (\mathcal{L}_1)^2$, and in general

$$\sigma_{-k}^F \sim \sum (\mathcal{L}_k)^{\alpha_k} (\mathcal{L}_{k-1})^{\alpha_{k-1}} \cdots (\mathcal{L}_1)^{\alpha_1}$$

with the summation taken over $0 \leq \alpha_i \leq k$, for every i and $\alpha_1 + 2\alpha_2 + \cdots + k\alpha_k = k$. The proof is complete. $\square$

REFERENCES

- [1] P. M. Alberti and R. Matthes, *Connes' trace formula and Dirac realization of Maxwell and Yang-Mills action*, arXiv:math-ph/991001.
- [2] T. P. Branson, *Conformally Covariant Equations on Differential Forms*, Comm. In Partial Differential Equations, **7**(4) (1982), 393–431.
- [3] T. P. Branson, *Differential operators canonically associated to a conformal structure*, Math. Scand., **57** (1985), 293–345.
- [4] A. Connes, *Noncommutative Geometry*, Academic Press, London and San Diego, 1994.
- [5] A. Connes, *Quantized calculus and applications*, XIth International Congress of Mathematical Physics (Paris, 1994), Internatl. Press, Cambridge, MA, 1995, pp. 15–36.
- [6] P. B. Gilkey, *Invariance Theory, the Heat Equation, and the Atiyah–Singer Index Theorem*, 2nd edition, CRC Press, Boca Raton, FL, 1995.
- [7] J. M. Gracia-Bondía, J. C. Várilly and H. Figueroa, *Elements of Noncommutative Geometry*, Birkhäuser Advanced Texts, Birkhäuser, Boston, 2001.
- [8] R. Graham, R. Jenne, L. Mason and G. Sparling, *Conformally invariant powers of the Laplacian, I: Existence*, J. London Math. Soc. (2) **46** (1992), 557–565.
- [9] A. Gray, *The volume of a small geodesic ball of a Riemannian manifold*, Michigan Math. J. **20** (1973), 329–344.
- [10] H. Kumano-go, *Pseudo-Differential Operators*, The MIT Press, Cambridge, Massachusetts, 1981.
- [11] F. Trèves, *Introduction to Pseudodifferential and Fourier Integral Operators I*, Plenum Press, New York, 1980.
- [12] W. J. Ugalde, *Differential forms canonically associated to even-dimensional compact conformal manifolds.*, to appear, Proceedings from “The 6th Conference on Clifford Algebras and their Applications in Mathematical Physics”, Tennessee Technological University, Cookeville, Tennessee, May 20–25, 2002.
- [13] F. W. Warner, *Foundations of Differential Manifolds and Lie Groups*, Springer-Verlag, New York, 1983.
- [14] M. Wodzicki, *Noncommutative residue. Chapter I: Fundamentals*, in *K-theory, Arithmetic and Geometry*, Yu. I. Manin, ed., Lecture Notes in Math. **1289**, Springer, Berlin, 1987, pp. 320–399.

William J. Ugalde
 Department of Mathematics
 14 MacLean Hall
 The University of Iowa
 Iowa City, Iowa, 52242
 E-mail: wugalde@math.uiowa.edu